

# The Evening Sky Map

FREE\* EACH MONTH FOR YOU TO EXPLORE, LEARN & ENJOY THE NIGHT SKY

## Sky Calendar – February 2016

Get Sky Calendar on Twitter  
<http://twitter.com/skymaps>

- 1 Last Quarter Moon at 3:28 UT.
  - 1 Moon near Mars (87° from Sun, morning sky) at 10h UT. Mag. +0.8.
  - 3 Moon near Saturn (60° from Sun, morning sky) at 19h UT. Mag. +0.5.
  - 6 Moon near Venus (30° from Sun, morning sky) at 7h UT. Mag. -4.0.
  - 6 Moon near Mercury (26° from Sun, morning sky) at 16h UT. Mag. 0.0. A good opportunity to view this elusive planet.
  - 7 Mercury at greatest elongation, 26° west of Sun (morning sky) at 1h UT. Mag. 0.0.
  - 8 New Moon at 14:39 UT. Start of lunation 1152.
  - 11 Moon at perigee (closest to Earth) at 3h UT (364,360 km; angular size 32.8').
  - 13 Mercury 4.0° E of Venus (25° and 29° from Sun, morning sky) at 3h UT. Mags. -0.1 and -4.0.
  - 15 First Quarter Moon at 7:46 UT.
  - 15 Moon near the Pleiades (evening sky) at 14h UT.
  - 16 Moon very near Aldebaran (evening sky) at 6h UT. Occultation visible from Hawaii, Philippines, Taiwan.
  - 20 Moon near Beehive cluster (evening sky) at 17h UT.
  - 22 Moon near Regulus (midnight sky) at 12h UT.
  - 22 Full Moon at 18:20 UT.
  - 24 Moon near Jupiter (165° from Sun, morning sky) at 3h UT. Mag. -2.5.
  - 26 Moon near Spica (morning sky) at 23h UT.
  - 27 Moon at apogee (farthest from Earth) at 3h UT (distance 405,383 km; angular size 29.5').
  - 29 Leap Day added to February so this year is 366 days long.
  - 29 Moon near Mars (morning sky) at 20h UT. Mag. +0.8.
- The Zodiacal Light** is caused by sunlight reflected off meteoric dust in the plane of the solar system. Choose a clear, moonless night, about 1–2 hours after sunset, and look for a large triangular-shaped glow extending up from the horizon (along the ecliptic). The best months to view the Zodiacal Light is when the ecliptic is almost vertical at the horizon: March and April (evening) and October–November (morning); times reversed for the southern hemisphere.

More sky events and links at <http://Skymaps.com/skycalendar/>  
 All times in Universal Time (UT). (USA Eastern Standard Time = UT - 5 hours.)



SAVE ON RECOMMENDED PRODUCTS • <http://Skymaps.com/store>


- STAR ATLASES & PLANISPHERES
  - STAR CHARTS & ASTRO POSTERS
  - BOOKS FOR SKY WATCHERS
  - TELESCOPES & BINOCULARS
- Help support the production and free distribution of The Evening Sky Map

## NORTHERN HEMISPHERE FEBRUARY 2016

SKY MAP SHOWS HOW THE NIGHT SKY LOOKS

EARLY FEB 8 PM  
 LATE FEB 7 PM

SKY MAP DRAWN FOR A LATITUDE OF 40° NORTH AND IS SUITABLE FOR LATITUDES UP TO 15° NORTH OR SOUTH OF THIS



- Symbols**
- Galaxy ☾
  - Double Star ●●
  - Variable Star ⊙
  - Diffuse Nebula □
  - Planetary Nebula ☆
  - Open Star Cluster ○
  - Global Star Cluster ⊕

**Star Magnitudes** -1 0 1 2 3 4

Copyright © 2000–2016 Kym Thalassoudis. All Rights Reserved.  
 \* TERMS OF USE: FREE FOR NON-COMMERCIAL EDUCATIONAL USE. ASTRONOMY EDUCATION GROUPS MAY FREELY DISTRIBUTE PRINTED HANDOUTS. FULL DETAILS AT <http://Skymaps.com/terms.html>

INSTRUCTIONS: THE SKY MAP SHOWS THE ENTIRE NIGHT SKY FROM HORIZON-TO-HORIZON AS IT APPEARS ON CERTAIN DATES AND TIMES. THE CENTER OF THE MAP IS THE PART OF THE SKY DIRECTLY OVERHEAD (ZENITH) AND THE OUTER RING IS THE PART OF THE SKY ON THE HORIZON. CELESTIAL OBJECTS ARE LOCATED BETWEEN THE ZENITH AND THE HORIZON. COMPASS DIRECTIONS ARE INDICATED ALONG THE HORIZON CIRCLE. (FOR EXAMPLE, NORTH.)

Use the Big Dipper (or Plough) located in the north-eastern sky to find Polaris, the North Star. In ancient times, the twins of Greek mythology, Castor & Pollux, were considered patron saints of seafarers. The ecliptic is almost vertical at the horizon: March and April (evening) and October–November (morning); times reversed for the southern hemisphere.

The mythical queen Cassiopeia is depicted in the sky by the thousand Cepheids & daughter (Andromeda). From northern latitudes, stars appear to rotate around the North Celestial Pole (NCP).

The Pleiades star cluster is more than twice the distance of the Hyades star cluster.

Magnificent Orion dominates the southern sky. The constellation represents a hunter with his two dogs nearby (Canis Major & Canis Minor).

The Big Dipper (or Plough) located in the north-eastern sky to find Polaris, the North Star. In ancient times, the twins of Greek mythology, Castor & Pollux, were considered patron saints of seafarers.

The ecliptic is almost vertical at the horizon: March and April (evening) and October–November (morning); times reversed for the southern hemisphere.

The Pleiades star cluster is more than twice the distance of the Hyades star cluster.

Magnificent Orion dominates the southern sky. The constellation represents a hunter with his two dogs nearby (Canis Major & Canis Minor).

The ecliptic is almost vertical at the horizon: March and April (evening) and October–November (morning); times reversed for the southern hemisphere.

The Pleiades star cluster is more than twice the distance of the Hyades star cluster.

Magnificent Orion dominates the southern sky. The constellation represents a hunter with his two dogs nearby (Canis Major & Canis Minor).

The ecliptic is almost vertical at the horizon: March and April (evening) and October–November (morning); times reversed for the southern hemisphere.

The Pleiades star cluster is more than twice the distance of the Hyades star cluster.

Magnificent Orion dominates the southern sky. The constellation represents a hunter with his two dogs nearby (Canis Major & Canis Minor).

The ecliptic is almost vertical at the horizon: March and April (evening) and October–November (morning); times reversed for the southern hemisphere.

The Pleiades star cluster is more than twice the distance of the Hyades star cluster.

Magnificent Orion dominates the southern sky. The constellation represents a hunter with his two dogs nearby (Canis Major & Canis Minor).

The ecliptic is almost vertical at the horizon: March and April (evening) and October–November (morning); times reversed for the southern hemisphere.

## About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. **Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars.** They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

## Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

## Astronomical Glossary

**Conjunction** – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

**Constellation** – A defined area of the sky containing a star pattern.

**Diffuse Nebula** – A cloud of gas illuminated by nearby stars.

**Double Star** – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

**Ecliptic** – The path of the Sun's center on the celestial sphere as seen from Earth.

**Elongation** – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

**Galaxy** – A mass of up to several billion stars held together by gravity.

**Globular Star Cluster** – A ball-shaped group of several thousand old stars.

**Light Year (ly)** – The distance a beam of light travels at 300,000 km/sec in one year.

**Magnitude** – The brightness of a celestial object as it appears in the sky.

**Open Star Cluster** – A group of tens or hundreds of relatively young stars.

**Opposition** – When a celestial body is opposite the Sun in the sky.

**Planetary Nebula** – The remnants of a shell of gas blown off by a star.

**Universal Time (UT)** – A time system used by astronomers. Also known as Greenwich Mean Time. USA Eastern Standard Time (for example, New York) is 5 hours behind UT.

**Variable Star** – A star that changes brightness over a period of time.

# NORTHERN HEMISPHERE FEBRUARY 2016 CELESTIAL OBJECTS Sky maps.com

## Easily Seen with the Naked Eye

Capella	Aur	•	The 6th brightest star. Appears yellowish in color. Spectroscopic binary. Dist=42 ly.
Sirius	CMa	•	The brightest star in the sky. Also known as the "Dog Star". Dist=8.6 ly.
Procyon	CMi	•	Greek name meaning "before the dog" - rises before Sirius (northern latitudes). Dist=11.4 ly.
δ Cephei	Cep	•	Cepheid prototype. Mag varies between 3.5 & 4.4 over 5.366 days. Mag 6 companion.
Deneb	Cyg	•	Brightest star in Cygnus. One of the greatest known supergiants. Dist=3,000 ly.
Castor	Gem	•	Multiple star system with 6 components. 3 stars visible in telescope. Dist=52 ly.
Pollux	Gem	•	With Castor, the twin sons of Leda in classical mythology. Dist=34 ly.
Regulus	Leo	•	Brightest star in Leo. A blue-white star with at least 1 companion. Dist=77 ly.
Rigel	Ori	•	The brightest star in Orion. Blue supergiant star with mag 7 companion. Dist=770 ly.
Betelgeuse	Ori	•	One of the largest red supergiant stars known. Diameter=300 times that of Sun. Dist=430 ly.
Algol	Per	•	Famous eclipsing binary star. Magnitude varies between 2.1 & 3.4 over 2.867 days.
Pleiades	Tau	•	The Seven Sisters. Spectacular cluster. Many more stars visible in binoculars. Dist=380 ly.
Hyades	Tau	•	Large V-shaped star cluster. Binoculars reveal many more stars. Dist=152 ly.
Aldebaran	Tau	•	Brightest star in Taurus. It is not associated with the Hyades star cluster. Dist=65 ly.
Polaris	UMi	•	The North Pole Star. A telescope reveals an unrelated mag 8 companion star. Dist=433 ly.

## Easily Seen with Binoculars

M31	And	•	The Andromeda Galaxy. Most distant object visible to naked eye. Dist=2.5 million ly.
M38	Aur	•	Stars appear arranged in "pi" or cross shape. Dist=4,300 ly.
M36	Aur	•	About half size of M38. Located in rich Milky Way star field. Dist=4,100 ly.
M37	Aur	•	Very fine star cluster. Discovered by Messier in 1764. Dist=4,400 ly.
M44	Cnc	•	Praesepe or Beehive Cluster. Visible to the naked eye. Dist=590±20 ly.
M41	CMa	•	First recorded observation by Aristotle in 325 BC as "cloudy spot". Dist=2,300 ly.
μ Cephei	Cep	•	Herschel's Garnet Star. One of the reddest stars. Mag 3.4 to 5.1 over 730 days.
Mira	Cet	•	Famous long period variable star. Mag varies between 3.0 & 10.1 over 332 days.
M39	Cyg	•	May be visible to the naked eye under good conditions. Dist=900 ly.
M35	Gem	•	Fine open cluster located near foot of the twin Castor. Dist=2,800 ly.
M48	Hya	•	12+ stars in 7x binoculars. Triangular asterism near centre. Dist=1,990 ly.
γ Leporis	Leo	•	Visible with binoculars. Gold & white stars. Mags 3.6 & 6.2. Dist=30 ly. Sep=96.3".
2232	Mon	•	A large scattered star cluster of 20 stars. Dist=1,300 ly.
2244	Mon	•	Surrounded by the rather faint Rosette Nebula. Dist=5,540 ly.
M50	Mon	•	Visible with binoculars. Telescope reveals individual stars. Dist=3,000 ly.
Cr 69	Ori	•	Lambda Orionis Cluster. Dist=1,630 ly.
M42	Ori	•	The Great Orion Nebula. Spectacular bright nebula. Best in telescope. Dist=1,300 light years.
Double Cluster	Per	•	Double Cluster in Perseus. NGC 869 & 884. Excellent in binoculars. Dist=7,300 ly.
M47	Pup	•	Bright star cluster. 15+ stars in 7x binoculars. Dist=1,500 ly.
M46	Pup	•	Dist=5,400 ly. Contains planetary NGC 2438 (Mag 11, d=65") - not associated.
Mizar & Alcor	UMa	•	Good eyesight or binoculars reveals 2 stars. Not a binary. Mizar has a mag 4 companion.

## Telescopic Objects

γ Andromedae	And	•	Attractive double star. Bright orange star with mag 5 blue companion. Sep=9.8".
γ Arietis	Ari	•	Impressive looking double blue-white star. Visible in a small telescope. Sep=7.8".
M67	Cnc	•	Contains 500+ stars mag 10 & fainter. One of the oldest clusters. Dist=2,350 ly.
M94	CVn	•	Compact nearly face-on spiral galaxy. Dist=15 million ly.
M51	CVn	•	Whirlpool Galaxy. First recognised to have spiral structure. Dist=25 million ly.
η Cassiopeiae	Cas	•	Yellow star mag 3.4 & orange star mag 7.5. Dist=19 ly. Orbit=480 years. Sep=12".
61 Cygni	Cyg	•	Attractive double star. Mags 5.2 & 6.1 orange dwarfs. Dist=11.4 ly. Sep=28.4".
θ Eridani	Eri	•	Striking blue-white double star. Mags 3.2 & 4.3. Visible in a small telescope. Sep=8.2".
γ Leonis	Leo	•	Superb pair of golden-yellow giant stars. Mags 2.2 & 3.5. Orbit=600 years. Sep=4.4".
β Monocerotis	Mon	•	Triple star. Mags 4.6, 5.0 & 5.4. Requires telescope to view arc-shape. Sep=7.3".
2264	Mon	•	Christmas Tree Cluster. Associated with the Cone Nebula. Dist=2,450 ly.
σ Orionis	Ori	•	Superb multiple star. 2 mag 7 stars one side, mag 9 star on other. Struve 761 triple in field.
k Puppis	Pup	•	Telescope easily shows two blue-white stars of almost equal brightness. Sep=9.9".
M1	Tau	•	Crab Nebula. Remnant from supernova which was visible in 1054. Dist=6,500 ly.
M33	Tri	•	Fine face-on spiral galaxy. Requires a large aperture telescope. Dist=2.3 million ly.
M81	UMa	•	Beautiful spiral galaxy visible with binoculars. Easy to see in a telescope.
M82	UMa	•	Close to M81 but much fainter and smaller.

## Calendario nocturno - Febrero 2016

- 1 Cuarto Creciente a las 3:28 TU.  
La Luna cerca de Marte (87 ° del Sol, cielo matutino) a las 10h TU. Mag. 0.8.
- 3 La Luna cerca de Saturno (60 ° del Sol, cielo de la mañana) de TU 19h. Mag. 0.5.
- 6 La Luna cerca de Venus (30 ° del Sol, cielo de la mañana) de TU 7h. Mag. -4.0.  
La Luna cerca de Mercurio (26 ° del Sol, cielo de la mañana) de TU 16h. Mag. 0.0.  
Una buena oportunidad para ver este planeta difícil de alcanzar.
- 7 Mercurio en su mayor elongación, 26 ° al oeste del Sol (cielo matutino) en TU 1h. Mag. 0.0.
- 8 Luna Nueva a las 14:39 TU. Comience de lunación 1152.
- 11 Luna en el perigeo (el más cercano a la Tierra) a 3h TU (364360 kilometros; tamaño angular de 32,8 ').
- 13 Mercurio 4.0 ° E de Venus (25 ° y 29 ° del Sol, cielo de la mañana) de TU 3h. Mags. -0,1 Y -4,0.
- 15 Luna Creciente a las 7:46 TU.  
La Luna cerca de las Pléyades (cielo nocturno) a las 14h TU.
- 16 La Luna muy cerca de Aldebaran (cielo nocturno) a 6h TU. Ocultación visible desde Hawaii, Filipinas, Taiwán.
- 20 La Luna cerca de la colmena cluster (cielo nocturno) a las 17h TU.
- 22 La Luna cerca de Regulus (cielo de medianoche) a las 12h TU.  
Luna llena a las 18:20 TU.
- 24 La Luna cerca de Júpiter (165 ° del Sol, cielo de la mañana) de TU 3h. Mag. -2.5.
- 26 La Luna cerca de Spica (cielo matutino) a las 23h TU.
- 27 Luna en apogeo (más alejada de la Tierra) a 3h TU (distancia 405.383 kilometros; tamaño angular de 29,5 ').
- 29 Día bisiesto añadido a febrero por lo que este año es de 366 días.  
La Luna cerca de Marte (cielo matutino) a las 20h TU. Mag. 0.8.

La luz zodiacal es causada por la luz solar reflejada por el polvo meteórico en el plano del sistema solar. Elija una clara noche sin luna, alrededor de 1-2 horas después de la puesta del sol, y busquer un gran resplandor de forma triangular que se extiende hasta el horizonte (a lo largo de la eclíptica). Los mejores meses para ver la luz zodiacal son es cuando la eclíptica está casi vertical en el horizonte: marzo y abril (de la noche) y octubre-noviembre (mañana); se invierte para el hemisferio sur.

Todas las horas Tiempo Universal (TU). ¡Un cielo despejado hasta el mes que viene!