

About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. **Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars.** They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

Astronomical Glossary

Conjunction – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

Constellation – A defined area of the sky containing a star pattern.

Diffuse Nebula – A cloud of gas illuminated by nearby stars.

Double Star – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

Elongation – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

Galaxy – A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) – The distance a beam of light travels at 300,000 km/sec in one year.

Magnitude – The brightness of a celestial object as it appears in the sky.

Open Star Cluster – A group of tens or hundreds of relatively young stars.

Opposition – When a celestial body is opposite the Sun in the sky.

Planetary Nebula – The remnants of a shell of gas blown off by a star.

Universal Time (UT) – A time system used by astronomers. Also known as Greenwich Mean Time. USA Eastern Standard Time (for example, New York) is 5 hours behind UT.

Variable Star - A star that changes brightness over a period of time.

NORTHERN HEMISPHERE MAY 2018

☆

M104

γ Virainis

Capella

Easily Seen with the Naked Eye

Arcturus	R00 ●	Orange, grant K star. Name means dear watcher . Dist=36.7 ty.
Procyon	CMi •	Greek name meaning "before the dog" - rises before Sirius (northern latitudes). Dist=11.4 ly.
δ Cephei	Cep ⊚	Cepheid prototype. Mag varies between 3.5 & 4.4 over 5.366 days. Mag 6 companion.
Deneb	Cyg •	Brightest star in Cygnus. One of the greatest known supergiants. Dist=1,400±200 ly.
Castor	Gem •	Multiple star system with 6 components. 3 stars visible in telescope. Dist=52 ly.
Pollux	Gem ●	With Castor, the twin sons of Leda in classical mythology. Dist=34 ly.
α Herculis	Her ⊚	Semi-regular variable. Magnitude varies between 3.1 & 3.9 over 90 days. Mag 5.4 companion.

• The 6th brightest star. Appears yellowish in color. Spectroscopic binary. Dist=42 ly.

Regulus Leo • Brightest star in Leo. A blue-white star with at least 1 companion. Dist=77 ly.

Vega Lyr • The 5th brightest star in the sky. A blue-white star. Dist=25.0 ly.

Antares Sco • Red, supergiant star. Name means "rival of Mars". Dist=135.9 ly.

Polaris UMi • The North Pole Star. A telescope reveals an unrelated mag 8 companion star. Dist=433 ly.

Spica Vir • Latin name means "ear of wheat" and shown held in Virgo's left hand. Dist=250 ly.

Easily Seen with Binoculars

		-	
μ Cephei	Сер	•	Herschel's Garnet Star. One of the reddest stars. Mag 3.4 to 5.1 over 730 days.
Mel 111	Com	0	Coma Berenices. 80 mag 5-6 stars in 5 deg. Dist=283 ly. Age=400 million years.
χ Cygni	Cyg	•	Long period pulsating red giant. Magnitude varies between 3.3 & 14.2 over 407 days.
M39	Cyg	0	May be visible to the naked eye under good conditions. Dist=900 ly.
ν Draconis	Dra	•	Wide pair of white stars. One of the finest binocular pairs in the sky. Dist=100 ly.
M13	Her	\oplus	Best globular in northern skies. Discovered by Halley in 1714. Dist=23,000 ly.
M92	Her	0	Fainter and smaller than M13. Use a telescope to resolve its stars.
R Hydrae	Hya	•	Long period variable. Mag varies between 3.0 & 11.0 over 390 days. Brilliant red.
ε Lyrae	Lyr	•	Famous Double Double. Binoculars show a double star. High power reveals each a double.
R Lyrae	Lyr	•	Semi-regular variable. Magnitude varies between 3.9 & 5.0 over 46.0 days.
M12	0ph	0	Close to the brighter M10. Dist=18,000 ly.
M10	0ph	\oplus	3 degrees from the fainter M12. Both may be glimpsed in binoculars. Dist=14,000 ly.

Cnc Praesepe or Beehive Cluster. Visible to the naked eye. Dist=590±20 ly.

• Easy to find in binoculars. Might be glimpsed with the naked eye.

IC 4665 Oph Carge, scattered open cluster. Visible with binoculars.

Oph Cattered open cluster. Visible with binoculars.

M4 Sco & A close globular. May just be visible without optical aid. Dist=7,000 ly.

M5 Ser & Fine globular star cluster. Telescope will reveal individual stars. Dist=25,000 ly.

Mizar & Alcor UMa • Good eyesight or binoculars reveals 2 stars. Not a binary. Mizar has a mag 4 companion.

Cr 399 Vul Coathanger asterism or "Brocchi's Cluster". Not a true star cluster. Dist=218 to 1,140 ly.

Telescopic Objects

reiescol	ppic Objects				
ε Boötis	Boo	•	Red giant star (mag 2.5) with a blue-green mag 4.9 companion. Sep=2.8". Difficult to split.		
M67	Cnc	\Diamond	Contains 500+ stars mag 10 & fainter. One of the oldest clusters. Dist=2,350 ly.		
M94	CVn	0	Compact nearly face-on spiral galaxy. Dist=15 million ly.		
η Cassiopeiae	Cas	•	Yellow star mag 3.4 & orange star mag 7.5. Dist=19 ly. Orbit=480 years. Sep=12".		
5128	Cen	0	Bisected by a wide obscuring lane. Strong radio source. Dist=14 million ly.		
M51	CVn	0	Whirlpool Galaxy. First recognised to have spiral structure. Dist=25 million ly.		
M64	Com	0	Black-Eye Galaxy. Discovered by J.E. Bode in 1775 - "a small, nebulous star".		
Albireo	Cyg	•	Beautiful double star. Contrasting colours of orange and blue-green. Sep=34.4".		
61 Cygni	Cyg	•	Attractive double star. Mags 5.2 & 6.1 orange dwarfs. Dist=11.4 ly. Sep=28.4".		
3242	Hya		Ghost of Jupiter. Bright blue disk. Mag 11 central star. Dist=2,600 ly.		
M83	Hya	0	Classic face-on spiral. Discovered in 1752 by Lacaille. In attractive star field.		
γ Leonis	Leo	•	Superb pair of golden-yellow giant stars. Mags 2.2 & 3.5. Orbit=600 years. Sep=4.4".		
β Lyrae	Lyr	•	Eclipsing binary. Mag varies between 3.3 & 4.3 over 12.940 days. Fainter mag 7.2 blue star.		
M57	Lyr		Ring Nebula. Magnificent object. Smoke-ring shape. Dist=4,100 ly.		
M16	Ser		Eagle Nebula. Requires a telescope of large aperture. Dist=8,150 ly.		
M81	UMa	0	Beautiful spiral galaxy visible with binoculars. Easy to see in a telescope.		
M82	UMa	0	Close to M81 but much fainter and smaller.		

Sombrero Galaxy. Almost edge-on spiral galaxy. Protruding central core.

Superb pair of mag 3.5 yellow-white stars. Orbit=169 years. At their closest in 2005.

♦ Dumbbell Nebula. Large, twin-lobed shape. Most spectacular planetary. Dist=975 ly.



Calendario de cielo - mayo de 2018

- 2 Luna cerca de Antares (cielo matutino) a las 11h UT.
- Venus 6.4 ° N de Aldebarán (28 ° del Sol, cielo vespertino) a las 18h UT. Mags. -3.9 y 0.9.
- 4 Luna cerca de Saturno (cielo matutino) a las 20h UT. Mag. 0.3.
- 6 Lluvia de meteoritos Eta Aquaridas. Más activa durante los 7 días cercanos a este día. Asociada con el cometa Halley. Meteoritos muy rápidos y brillantes, picos máximos de hasta 30 por hora. Favorece a los observadores del cielo en los trópicos observando unas horas antes del amanecer. Condiciones de visión desfavorables este año debido a la brillante luz de la luna.
 - Luna en el apogeo (más alejado de la Tierra) a la 1h UT (distancia 404,457 km, tamaño angular 29,5 ').
 - Luna cerca de Marte (cielo matutino) a las 7h UT. Mag. -0.5.
- 8 Luna en cuarto menguante a las 2:10 UT.
- 9 Júpiter en la oposición a las 0h UT. Mejor momento para observar el planeta más grande del sistema solar. Mag. -2.5.
- 13 Luna cerca de Mercurio (23 ° del Sol, cielo matutino) a las 19h UT. Mag. -0.2.
- Luna Nueva a las 11:49 UT. Inicio de la lunación 1180.
- 16 Luna cerca de Aldebarán (15 ° del Sol, cielo vespertino) a las 13h UT.
- Luna cerca de Venus (31 ° del Sol, cielo vespertino) a las 19h UT. Mag. -4.0.
 Luna en perigeo (más cercana a la Tierra) a las 21h UT (363,776 km, tamaño angular 32,8 ').
- 19 Luna cerca de Pollux (cielo vespertino) a las 13h UT.
- 20 Luna cerca del cúmulo de Beehive M44 (cielo vespertino) a las 12h UT.)
- Venus 0.7 ° N del grupo M35 (32 ° del Sol, cielo vespertino) a las 10h UT. Mags. -4.0 y 5.1.
- Luna cerca de Regulus (cielo vespertino) a las 2h UT.
 - Luna en cuarto creciente a las 3:50 UT.
- Luna cerca de Spica (cielo vespertino) a las 2h UT.
- 27 Luna cerca de Júpiter (cielo vespertino) a las 20h UT. Mag. -2.5.
- 29 Luna Llena a las 14:20 UT.
 - Luna cerca de Antares (cielo de medianoche) a las 19h UT.

Todos los tiempos Universal Time (UT). ¡Cielos despejados hasta el próximo mes!